

Intoxication by *Cortinarius orellanus*: Detection and assay of orellanine in biological fluids and renal biopsies

S. Rapior,¹ N. Delpech,² C. Andary¹ & G. Huchard³

¹Laboratoire de Botanique, Phytochimie et Mycologie, Faculté de Pharmacie, Avenue Charles Flahault, 34060 Montpellier Cedex 1, France; ²Laboratoire de Biochimie-Immunochimie, Centre Hospitalier Général, B.P. 4045, 66042 Perpignan Cedex, France; ³Service de Néphrologie-Hémodialyse, Centre Hospitalier Général, B.P. 4045, 66042 Perpignan Cedex, France

Accepted 15 April 1989

Key words: *Cortinarius orellanus*, nephrotoxicity, orellanine, assay, treatment

Abstract

A woman suffering from acute tubulo-interstitial nephritis was admitted to the hospital ten days after deliberate intoxication by ingestion of *Cortinarius orellanus*. Orellanine, the main toxin responsible for orellanine poisoning, was detected in biological fluids and renal biopsies. It was assayed by direct spectrofluorimetry on two-dimensional thin-layer chromatograms after specific photodecomposition into orelline. The orellanine concentration was 6.12 mg/l in the plasma (10 days after ingestion). Orellanine levels in renal biopsies were 7 µg per 25 mm³ of the first biopsy (13 days after ingestion) and 24 µg per 8 mm³ of the second biopsy (6 months later).

Introduction

Many cases of orellanine poisoning were described after several fatal intoxications in Europe. They were attributed to *Cortinarius orellanus* (Fr.) Fr. [5, 8, 10, 17, 18, 19, 32, 33] and *Cortinarius speciosissimus* Kühn & Romagn. [6, 9, 13, 14, 20, 22, 29, 30, 31, 38, 39]. Following the isolation of orellanine by Grzymala [11], Antkowiak & Gessner [4] reported a tetra-hydroxy bipyridine, N,N-dioxide structure for this molecule. This was confirmed by synthesis [7, 34].

We recently described [2, 26] the first case of deliberate intoxication by *C. orellanus*. This report was confirmed by assaying orellanine in the

plasma and two renal biopsies of a patient after specific photodecomposition into a non-toxic metabolite called orelline [1, 3]. The originality of our work is primarily the reporting of the experimental conditions of detection and assay of orellanine in biological tissues [2, 26].

Materials and methods

Case report

A 31-year-old female psychiatric patient was brought to hospital ten days after deliberate ingestion of two fruit-bodies of *C. orellanus* (Novem-

* Taken in part from the doctorate thesis of S. Rapior.

ber, 1987). She experienced severe thirst, nausea and abdominal pains, and vomiting respectively eight and nine days after ingestion. On day 10 after ingestion, the woman had a high plasma creatinine level (1100 $\mu\text{mol/l}$) and a severe renal failure. She was afebrile and anicteric (SGPT: 5 UI/l; SGOT: 12 UI/l; LDH: 370 UI/l). Haemodialysis with femoral catheters (500 ml/mn for 3 hours) and perfusion with furosemide (Lasilix), diltiazem (Tildiem), dopamine and vitamin C were started immediately on the night of admission. The treatment was repeated on the following five days (except for Tildiem up to nine days). On days 11 to 13 and 17 to 19 after ingestion (Table 1), haemodialysis was performed with Amberlite XAD-7 resin (ref. DHP-1, Kurakay Co., Japan and ref. Hemo France, France). On day 16, the patient's plasma was exchanged for 3% albumine in saline combined with a haemoperfusion through Amberlite XAD resin for 90 mn. Parenteral 30% glucose solution (60 ml), isotonic bicarbonate solution (2×500 ml) and Calciparine were administrated continuously during perfusion. On days 11 to 19 after ingestion, the patient was perfused with an amino acid mixture (Protinutril, Roger Bellon

Laboratories, France) (500 ml/24 h) and given another amino acid mixture in the form of four tablets three times a day (Ketosteril, Fresenius Laboratories, FGR). Renal biopsies were performed on days 13 and 180 after ingestion.

Histologic failures were those of a marked tubular necrosis with scattered infiltration of lymphocytes and interstitial oedema and fibrosis. Plasma creatinine decreased to 350 $\mu\text{mol/l}$ and clearance creatinine remained at about 10 to 20 ml/mn for the next days. On day 27 after ingestion, the woman returned home with a sufficient renal function to maintain life without periodic dialysis (plasma creatinine: 383 $\mu\text{mol/l}$; uraemia: 24 mmol/l; red corpuscles: $2.81 \cdot 10^6/\text{mm}^3$; hemoglobin: 8 g/100 ml; hematocrit: 24%). Four months after ingestion, plasma creatinine (240 $\mu\text{mol/l}$) and uraemia (11.1 mmol/l) decreased and urinary excretion of proteins remained at about 0.44 g/24 h. Six months after ingestion, blood analysis gave the following results: plasma creatinine: 209 $\mu\text{mol/l}$; uraemia: 11.5 mmol/l; red corpuscles: $3.94 \cdot 10^6/\text{mm}^3$; hemoglobin: 10.9 g/100 ml; hematocrit: 32.5%; urinary creatinine: 4.1 mmol/l; clearance: 0.4541 ml/s.

Table 1. Chronology of biological samples.

Days ^a	Plasma ^b	Urine	Renal biopsy	Haemodialysis fluid	Haemodialysis resin	Plasmapheresis resin
10	P10B +	-	-	HF10	HR10 +	-
11	P11B +	-	-	HF11	HR11 +	-
	P11A +	-	-			
12	P12B +	-	-	HF12 +	HR12 +	-
	P12A +	-	-			
13	P13A +	-	RB13 +	HF13 +	HR13 +	-
14	-	-	-	-	-	-
15	-	U15 +	-	-	-	-
16	-	U16 +	-	-	-	PR16 +
17	-	-	-	HF17	HR17	-
18	-	-	-	HF18	HR18	-
19	-	-	-	HF19	HR19	-
180	-	-	RB180 +	-	-	-

^a Days after ingestion.

^b A: after haemodialysis, B: before haemodialysis.

+ Analyzed samples.

- No sample.

Purification of orellanine in biological fluids and renal biopsies

Orellanine and its photodecomposition metabolites were purified by passage through a 22 × 200 mm column containing Amberlite XAD-4 resin (Rohm & Haas, ref. 2-6007) adjusted to pH 2 with hydrochloric acid. Elution was carried out by successive passages of following solvents: water pH 2 with hydrochloric acid, methanol-water (1:1) and methanol. The fractions were evaporated *in vacuo* to known volumes (about 0.5 ml).

Plasma samples (P10B to P13A) were loaded directly on the column. Urine samples (U15 and U16) and haemodialysis fluid samples (HF12 and HF13) were evaporated in a rotary vacuum evaporator to 5 ml before Amberlite XAD-4 resin treatment. Haemodialysis resin samples HR10 to HR13 and plasmapheresis resin sample PR16 were washed with water (30%, w/v). Orellanine was extracted by maceration of washed resin in methanol (12%, w/v) several times for periods of 15 mn under magnetic agitation at room temperature. The methanolic extracts were evaporated *in vacuo* to 2 ml. Tissue biopsies (RB13 and RB180) were homogenized with 1 ml of methanol-water (1:1) in a Potter homogenizer and then sonicated for 20 mn at room temperature. The hydromethanolic extract was centrifuged at 3000 g for 30 mn. The supernatant was evaporated *in vacuo* to 0.2 ml. Because of the instability of orellanine in light, this study was carried out in the dark [1, 15, 25].

Detection and assay of orellanine

Orellanine was revealed by two-dimensional tlc on cellulose (Merck, ref 5716) developed in the following solvent: *n*-butanol-hydrochloric acid-chloroform-water (40:20:15:3.8). The assay used the specific photodecomposition of orellanine into orelline under UV light (366 nm) for 90 mn through a quartz cell in a liquid environment. Fluorescence intensity of orelline was determined with a Farrand UV-Vis-2 chromatographic

analyzer (Optical, New York, USA) by direct spectrofluorimetry on two-dimensional chromatograms (10 ng detection limit on chromatograms). We compared 5 or 10 µl of each extract, previously exposed to UV light, with an orelline solution prepared with an orellanine standard solution (0.002% in methanol-water 1:1), irradiated with UV light in the same conditions (Fig. 1).

Results and discussion

Many cases of orellanine poisoning have been described in various European countries since 1952. Two cases histories reported by Marichal [18, 19] and Brousse [5] illustrated such poisoning in France. In September 1987, the failure to identify *Cortinarius* mushrooms led a collective poisoning by *C. orellanoides* of 26 soldiers in survival training [23].


The primary originality of our work is firstly the reporting of the first case of deliberate orellanine poisoning with accurate knowledge of the name of the *Cortinarius* species, the amount of mushroom eaten and the date of ingestion. This is very important because the clinical and histological aspects of orellanine poisoning are notable by virtue of the long incubation period (10 days). They were characterized in particular by a predominantly renal involvement (plasma creatinine: 1100 µmol/l) with tubulo-interstitial failures, infiltration of lymphocytes and subsequent development of fibrotic tissue. These findings are the same as those described in animal tests [16, 24]. No glomerular damage such as that reported in the literature [5, 10, 13, 17, 18, 19] was noted. Liver function tests were normal [5, 6, 19, 22].

The tubular damage with oliguric nephritis necessitated symptomatic early treatment with haemodialysis-haemoperfusion combined with plasmapheresis and perfusion with furosemide, dopamine and vitamin C for five days. The patient was also treated with amino acid perfusion and per os mixture with good results with regard to kidney preservation. According to Toback [35], amino acid perfusion acts directly on the kidneys to increase the synthesis of phospholipids

Orellanine →

RB13 0.1 0.2 0.3 μg
Orelline

a

RB13 0.2 0.3 0.4 μg
(+UV) Orelline

b

Fig. 1. Two-dimensional TLC on cellulose for orellanine assay in biological tissues. a: before photodecomposition; b: after photodecomposition; I: first development; II: second development; RB13: renal biopsy on day 13 after ingestion of *C. orellanus*.

containing choline for new membranes in regenerating cells and also to decrease the level of renal functional insufficiency after tubular necrosis. The vasodilatation effects of parenteral Tildiem have been used as theoretical protective treatment in acute renal failures [36, 37]. With respect to restoration of renal function, treatment by amino acids combined with Tildiem, haemodialysis and plasmapheresis-haemoperfusion combination might have been beneficial for the patient. After that, plasma creatinine and uraemia declined from 383 to 209 $\mu\text{mol/l}$ and from 24 to 11.5 mmol/l respectively for the six following months. Renal biopsy RB180 (six months after ingestion) showed tubular recovery but with intensive, evolutive interstitial fibrosis.

We were also able to detect and assay orellanine, the main toxin responsible for orellanine poisoning. The ability of Amberlite XAD resin (type 2, 4 or 7) to remove e.g. psychotropic drugs [12, 21] and phenolic type compounds [2, 26, 27, 28] from aqueous solutions led us to purify orellanine from biological fluids and renal biopsies with exclusion chromatography carried out on Amberlite XAD-4 resin. On account of its specific macroreticular structure and its very high surface area, Amberlite XAD-4 resin has a greater adsorption capacity for compounds with low molecular weights ($\text{MW} < 500$) i.e. orellanine than Amberlite XAD-2 or XAD-7 resins. According to Rosler & Goodwin [28], the initial acidification on the resin and the wash water-HCl may indirectly improve the capacity of the resin. Large molecules (i.e. creatinine) were excluded from the pore structure of the gel with water pH2 and passed straight through the column. On the other hand, small molecules (i.e. orellanine and its compounds of photodecomposition) were eluted with methanol-water (1:1) and methanol.

Orellanine was only partially separated by the mobile phase when a one-dimensional thin-layer chromatographic technique was used. We therefore employed a two-dimensional procedure in the same solvent for the present investigation. The identity of orellanine was confirmed by its specific photodecomposition into orelline under UV light. The identity of orellanine was confirmed by its

fluorescence spectrum in the chromatogram spot (excitation wavelength: 399 nm; emission wavelength: 447 nm).

This procedure led us to detect and assay orellanine (6.12 mg/l = 0.02 mmol/l) in plasma P10B (obtained before first haemodialysis). We did not detect orellanine and its photodecomposition products in other plasma or in urine, haemodialysis fluid or haemodialysis and plasmapheresis resin samples. On the other hand, we detected high amounts of orellanine and orelline and small amounts of its photodecomposition compounds in renal biopsy RB13 (13 days after ingestion). Orellanine level was 7 μg (= 0.02 μmol) per 25 mm³ of first renal biopsy. Renal biopsy RB180 (six months after ingestion) contained mainly orelline; its amount was equivalent to 24 μg (= 0.09 μmol) of orellanine per 8 mm³ of the second renal biopsy. Haemodialysis had thus cleared circulating toxin from the blood but had not been able to remove toxin bound to renal cells.

The characterization of orellanine in plasma obtained ten days after ingestion and the detection of orelline in renal biopsy six months later proved a particularly slow release of orellanine and its photodecomposition compounds from blood and renal cells. These metabolites may play a significant role in orellanine poisoning and might form covalent bonds with glutathione and with numerous proteins causing kidney disturbance. These findings support our hypothesis of the phototoxicity of orellanine [1]. The originality of our work is primarily related to the experimental conditions which made it possible to detect and assay orellanine in complex biological fluids and renal biopsies for the first time. Using this easy analytical procedure, it is now possible to assay this toxin anywhere in the body and to make an accurate diagnosis of orellanine intoxication.

References

1. Andary C, Rapior S, Fruchier A, Privat G. Cortinaires de la section *Orellani*: photodécomposition et hypothèse de la phototoxicité de l'orellanine. *Cryptogamie Mycol* 1986; 7: 189-200.

2. Andary C, Rapior S, Delpech N, Huchard G. Deliberate intoxication by ingestion of *Cortinarius orellanus*. Lancet 1989; 1: 213.
3. Antkowiak WZ, Gessner WP. Isolation and Characteristics of Toxic Components of *Cortinarius orellanus* Fries. Bull Acad Pol Sci Sér Sci Chimiques 1975; 23: 729-33.
4. Antkowiak WZ, Gessner WP. The structures of orellanine and orelline. Tetrahedron Lett 1979; 21: 1931-4.
5. Brousse A, Hervé JP, Leguy P, Cledes J, Leroy JP. L'intoxication par champignons de type *Cortinarius orellanus*. Une cause rare d'insuffisance rénale. Nouv Press Méd 1981; 10: 1940.
6. Busnach G, Dal Col A, Perrino ML, Surian M, Rovati C, Barbiano di Belgioioso G, Minetti L. Plasma exchange in acute renal failure by *Cortinarius speciosissimus*. Int J Artif Organs 1983; 6: 73-4.
7. Dehmlow EV, Schulz HJ. Synthesis of orellanine, the lethal poison of a toadstool. Tetrahedron Lett 1985; 26: 4903-6.
8. Färber D, Feldmeier S. Die *Orellanus*-Pilzvergiftung im Kindesalter. Anästhes Prax 1977; 13: 87-92.
9. Fauchald P, Westlie L. Poisoning by *Cortinarius speciosissimus*. Tidsskr Nor Laegeforen 1982; 102: 15-16.
10. Favre H, Leski M, Christeler P, Vollenweider E, Chatelangat F. Le *Cortinarius orellanus*: un champignon toxique provoquant une insuffisance rénale aiguë retardée. Schweiz med Wschr 1976; 106: 1097-102.
11. Grzymala S. L'isolement de l'orellanine, poison de *Cortinarius orellanus* Fries et l'étude de ses effets anatomopathologiques. Bull Soc Mycol France 1962; 78: 394-404.
12. Heath A, Delin K, Edén E, Martensson E, Selander D, Wickström I, Ahlmén J. Hemoperfusion with Amberlite Resin in the Treatment of Self-Poisoning. Acta Med Scand 1980; 207: 455-60.
13. Holmdahl J, Ahlmén J, Svalander C, Eriksson J, Bucht H. Renal damage after intoxication with *Cortinarius* mushrooms. Toxicological aspects. Ed An Kovatsis, Technika Studio (Greece) 1980: 155-63.
14. Hulmi S, Sipponen P, Forsström J, Vilska J. Mushroom poisoning caused by *Cortinarius speciosissimus*. A report of four cases. Duodecim 1974; 14-15: 1044-50.
15. Kurnsteiner H, Moser M. Isolation of a lethal toxin from *Cortinarius orellanus* Fr. Mycopathol 1981; 74: 65-72.
16. Lahtiperä S, Naukkarinen A, Collan Y. Mushroom Poisoning due to *Cortinarius speciosissimus*: Electron Microscope Study in Rats. Arch Toxicol 1986; Suppl 9: 315-19.
17. Leski M, Favre H, Chatelangat F, Vollenweider E, Baczkó AM. Insuffisance rénale aiguë provoquée par l'ingestion d'un champignon: *Cortinarius orellanus*. J Urol 1976; 82: 976-81.
18. Marichal JF, Carbiener R, Wiederkehr JL, Triby F. Insuffisance rénale chronique après intoxication par champignons de type *Cortinarius orellanus*. Coll Méd Lég Toxicol Méd n°106. Champignons toxiques. Compte-rendu de la Réunion du Groupement Français des Centres Anti-Poisons, Pont-à-Mousson 26 mai 1977; 161-5.
19. Marichal JF, Triby F, Wiederkehr JL, Carbiener R. Insuffisance rénale chronique après intoxication par champignons de type *Cortinarius orellanus* Fries. Nouv Press Méd 1977; 6: 2973-5.
20. Morild I, Giertsen JC, Christensen J. Mushroom poisoning as a cause of 'sudden death'. Tidsskr Nor Laegeforen 1983; 103: 1529-52.
21. Mulec H, Vikgren P, Nyqvist A. Hemoperfusion (HP) with Amberlite XAD-4 resin for treatment of intoxications in a county hospital. Artif Organs 1981; 5: 207.
22. Nolte S, Hufschmidt C, Steinhauer H, Rohrbach R, Künzer W. Terminale Niereninsuffizienz durch interstitielle Nephritis nach Pilzvergiftung durch *Cortinarius speciosissimus*. Monatsschr Kinderheilkd 1987; 135: 280-1.
23. Pats B, Durand G, Bousser J, Beauche A, Nicand E, Galliez M, Thomas R, Giudicelli CP. Toxicologie automnale lors d'un exercice de survie - A propos d'une intoxication collective par champignons de type *Cortinarius orellanoides*. Médecine et armées 1988; 16: 231-3.
24. Prast H, Pfäller W. Toxic properties of the mushroom *Cortinarius orellanus* (Fries). II. Impairment of renal function in rats. Arch Toxicol 1988; 62: 89-96.
25. Rapior S. Isolement et structure des orellanines chez *Cortinarius orellanus*. DEA Spécialité Chimie Thérapeutique. Faculté de Pharmacie, Université de Montpellier I 1983: 84 p.
26. Rapior S. Contribution à l'étude de *Cortinarius orellanus* (Fr.). Fr.: chimiotaxonomie, culture *in vitro*, toxicité. Thèse d'Etat Sciences Pharmaceutiques, Université de Montpellier I 1988: 254 p.
27. Rapior S, Andary C, Mousain D. *Cortinarius* section *Orellani*: Isolation and Culture of *Cortinarius orellanus*. Trans Br mycol Soc 1987; 89: 41-4.
28. Rosler KH, Goodwin RS. A general use of Amberlite XAD-2 resin for the purification of flavonoids from aqueous fractions. J Nat Prod 1984; 47: 188.
29. Schumacher T, Hoiland K. Toxic *Cortinarius* spp and *Cortinarius* poisoning. Naturen 1978; 3: 135-40.
30. Schumacher T, Hoiland K. Mushroom Poisoning Caused by Species of the Genus *Cortinarius* Fries. Arch Toxicol 1983; 53: 87-106.
31. Short AIK, Watling R, MacDonald MK, Robson JS. Poisoning by *Cortinarius speciosissimus*. Lancet 1980; 942-4.
32. Skirgiello A. Erfahrungen mit *Dermocybe orellana* (Fr.) in Polen - A: *Cortinarius (Dermocybe) orellanus* Fr. non Qué. - Cause d'intoxications fongiques en Pologne en 1952-1956. Zeitschr f Pilzk 1957; 23: 138-9.
33. Soresina P, Leoni G. Il *Cortinarius orellanus* ed i suoi effetti tossici. Mic Ital 1978; 3: 9-15.
34. Tiecco M, Tingoli M, Testaferri L, Chianelli D, Wenkert

- E. Total synthesis of orellanine, the lethal toxin of *Cortinarius orellanus* Fries mushroom. *Tetrahedron* 1986; 42: 1575-85.
35. Toback FG. Amino acid enhancement of renal regeneration after acute tubular necrosis. *Kidney Int* 1977; 12: 193-8.
36. Wagner K, Neumayer HH. Prevention of delayed graft function in cadaver kidney transplants by diltiazem. *Lancet* 1985; 1355-6.
37. Wagner K, Albrecht S, Neumayer HH. Prevention of Delayed Graft Function in Cadaveric Kidney Transplantation by a Calcium Antagonist. Preliminary Results of Two Prospective Randomized Trials. *Transplantation Proceedings* 1986; 18: 510-5.
38. Watling R. Orellanin-Poisoning in Britain. *Bull Br Mycol Soc* 1980; 14: 63.
39. Watling R. *Cortinarius speciosissimus*. The cause of renal failure in two young men. *Mycopathol* 1982; 79: 71-8.

